

Facial Nerve Regeneration

Arch Otolaryngol Head Neck Surg. 1998 Apr;124(4):383-9.

Effect of pulsed electromagnetic stimulation on facial nerve regeneration.

[Byers JM](#), [Clark KF](#), [Thompson GC](#).

Department of Otorhinolaryngology, University of Oklahoma Health Sciences Center, Oklahoma City, USA.

OBJECTIVE: To determine if exposure to electromagnetic fields influences regeneration of the transected facial nerve in the rat. **DESIGN AND METHODS:** The left facial nerve was transected in the tympanic section of the fallopian canal in 24 rats randomly assigned to 2 groups. The cut ends of the facial nerve were reapproximated without sutures within the fallopian canal to maximize the potential for regeneration. Rats in the experimental group (n= 12) were then exposed to pulsed electromagnetic stimulation (0.4 millitesla at 120 Hz) for 4 hours per day, 5 days per week, for 8 weeks. Rats in the control group (n=12) were handled in an identical manner without pulsed electromagnetic stimulation. Four other rats were given sham operations in which all surgical procedures were carried out except for the actual nerve transection. Two of these rats were placed in each group. Nerve regeneration was evaluated using electroneurography (compound action potentials), force of whisker and eyelid movements, and voluntary facial movements before and at 2-week intervals after transection. Histological evaluation was performed at 10 weeks after transection. Each dependent variable was analyzed using a 2-way analysis of variance with 1 between variable (groups) and 1 within repeated measures variable (days after transection). **RESULTS:** Statistical analysis indicated that N1 (the negative deflection of depolarization phase of the muscle and/or nerve fibers) area, N1 amplitude, and N1 duration, as well as absolute amplitude of the compound action potentials, were all significantly greater 2 weeks after transection in the experimental than in the control group of rats. The force of eye and whisker movements after electrical stimulation was statistically greater in the experimental group of rats 4 weeks after transection. Voluntary eye movements in the experimental group were significantly better at 5 and 10 weeks, while whisker movements were better at 3 and 10 weeks. There was no statistical difference between the 2 groups for any histological variable. **CONCLUSION:** Results of this study indicate that pulsed electromagnetic stimulation enhances early regeneration of the transected facial nerve in rats.

Zh Nevropatol Psikhiatr Im S S Korsakova. 1997;97(3):32-5.

[Magnetic and electromagnetic stimulation in the combined treatment of patients with paralysis of the mimetic musculature]

[Article in Russian]

Korotkikh NG, Korzh GM.

The original method of treatment of patients with postoperative paresis of mimic muscles was offered. The method included the use of two therapeutic factors: magnetotherapy and electromagnetostimulation of peripheral branches of facial nerve. The principles of combined influence of both components were described in detail. Data on functional control of treatment results confirm the reparation of peripheral branches of facial nerve.