

## Facial Pain

[Zhongguo Zhen Jiu](#). 2008 Apr;28(4):265-6.

### **[Observation on therapeutic effect of acupuncture combined with He-Ne laser radiation on facial paralysis]**

[Article in Chinese]

[Hou YL](#), [Li ZC](#), [Ouvang Q](#), [Li X](#), [Li H](#), [Zhao M](#).

Section of Physiotherapy, General Hospital of Beijing Military Area, Beijing 100700, China.

**OBJECTIVE:** To compare therapeutic effects of acupuncture combined with He-Ne laser radiation and western medicine on facial paralysis. **METHODS:** Ninety-seven cases were randomly divided into an acupuncture with laser group (n=55) and a medication group (n=42). The acupuncture with laser group were treated by acupuncture at main points, Dicang (ST 4), Jiache (ST 6), Sibai (ST 2), Qianzheng, Xiaguan (ST 7), Yingxiang (LI 20), Yangbai (GB 14), Yuyao (EX-HN 4), Yifeng (TE 17), Hegu (LI 4), in combination with He-Ne laser radiation; the medication group were treated with routine western medicine including intravenous dripping of low molecular dextran, compound Danshen and ATP injections, intramuscular injection of VitB1 and VitB12, and oral administration of oryzanol and prednisone. Seven days constituted one course. After two courses their therapeutic effects were observed. **RESULTS:** The cured rate was 81.8% in the acupuncture with laser group and 45.20% in the medication group with a significant difference between the two groups (P<0.01). **CONCLUSION:** The therapeutic effect of acupuncture combined with He-Ne laser radiation on facial paralysis is better than that of routine medication.

### **HELIUM-NEON LASERTHERAPY IN TREATMENT OF FACIAL NERVE NEUROPATHY**

A. Scherbonosova, V.V. Skupchenko  
Medical university, Far Eastern Medical Center, Khabarovsk, Russia

Facial nerve damage is the result of different factors influence and it appears at the background of ischemia anoxia. Elimination of a local pathologically fixed ergothromages ischemia condition with the help of helium-neon laser therapy (FTNLT)

has been conducted taking into consideration initial vegetative pattern of patients. It allowed to adjust treatment methods based on laser effect individually for every patient. Mimic muscles function restoration in the course of treatment had begun after 5 treatment sessions with HNLT and matched vegetative status normalization as well as reofaciogram, ultrasound Dopplerography of temporal and ophthalmic arteries and general conjunctive index. Thus, HNLT is a gentle corrector of vegetative homeostasis and sanogenic mechanisms. It allows to synchronize local and cerebral hemodynamics rhythms and trigger reparative regeneration of the facial nerve.

### **[Laser therapy and cryomassage in rehabilitation of patients with facial nerve neuropathy]**

[Article in Russian]

**Maslovskaja SG, Gusarova SA, Gorbunov FE, Strel'tsova EN.**

Cryomassage and its combination with low-intensity infra-red laser radiation have been introduced as a novel treatment of facial nerve neuropathy (FNN) in 32 patients. Electrophysiological investigations (facial thermography, classical electrodiagnosis, electromyography of the mimic muscles) and clinical data including those of long-term follow-up show that neither cryomassage nor infra-red laser radiation studied promote transformation of facial tissues in FNN patients. Use of the above factors is effective in a preclinical stage of forming contracture of the mimic muscles. Special techniques of application of local hypothermia and laser radiation can be used in multimodality treatment of both the established contracture and sluggish paresis of the facial muscles.

[Masui](#). 1992 Nov;41(11):1809-13.

### **[Low reactive-level laser therapy near the stellate ganglion for postherpetic facial neuralgia]**

[Article in Japanese]

[Ohtsuka H](#), [Kemmonsu O](#), [Dozaki S](#), [Imai M](#).

Department of Anesthesiology, Hokkaido University School of Medicine, Sapporo.

Low reactive-level laser therapy near the stellate ganglion was given for a 68-year-old female with postherpetic neuralgia, suffering from burning pain in the right forehead for 11 years. Stellate ganglion block and supraorbital nerve block with oral medication were not effective to relieve this pain. The laser irradiation induced warm sensation in her face followed by an excellent pain relief. Thermograms illustrated a remarkable increase from 30.6 degrees C to 31.5 degrees C in temperature of her right face. The irradiation near the right carotid artery also had the similar effect. The results imply that the irradiation with

low reactive-level laser of the stellate ganglion and/or the carotid artery increases a facial blood flow and relieves facial neuralgia.